Displaced lepton flavour violating signatures of right-handed sneutrinos in $U(1)'$ supersymmetric models

Priyotosh Bandyopadhyay*
Dipartimento di Matematica e Fisica "Ennio De Giorgi", Università del Salento and INFN-Lecce, Via Arnesano, 73100 Lecce, Italy†

We consider a $U(1)'$ extended supersymmetric model with a right handed neutrino superfield which can generate light neutrino mass by Type I seesaw mechanism. The lighter part of superpartner of the right-handed neutrino cloud be the candidate dark matter. Right-handed snuetrinos can come from the decay of $\tilde{Z}'$, superpartner of the extra gauge boson Z'. Left-right handed snuetrino mixing in this case can give rise to displaced lepton flavour violating signatures. Lighter right-handed snutrinos along with a wino-like chargino NLSP (next to lightest supersymmetric particle) creates an interesting decay topology. We investigate such displaced multi-leptonic final states with lepton flavour violation from the third generation supersymmetric cascade decays at the LHC.

arXiv:1511.03842v1 [hep-ph] 12 Nov 2015

The recent discovery of a Higgs boson around 125 GeV has established the role of at least one scalar in the electro-weak symmetry breaking [1]. However, experimental evidences for tiny neutrino masses and indirect evidence of dark matter candidate, and a theoretical requirement for naturalness of the electroweak scale needs a theory beyond Standard Model(SM). A supersymmetric seesaw model [2] with an additional $U(1)'$ [3] gauge symmetry can address these issues consistently.

In supersymmetric theories with R-parity, the lightest supersymmetric particle (LSP) is stable and thus a neutral LSP, typically a linear combination of neutral gauginos and Higgsinos, becomes a good thermal dark matter (DM) candidate if supersymmetry (SUSY) is broken around the TeV scale [4]. Similarly $U(1)'$ breaking can also be indued via quantum effects at $\mathcal{O}$(TeV) [5] which then generates the seesaw scale around TeV testable at LHC. The additional $U(1)'$ if comes from any grand unified gauge group which requires the presence of right-handed neutrinos (RHNs) for the anomaly-free condition, and cloud be at TeV-scale.

A right-handed sneutrino (RHsN) can be the LSP and a good dark matter candidate whose thermal relic density is in the right range if the $U(1)'$ gaugino $\tilde{Z}'$, the superpartner of the $U(1)'$ gauge boson Z', is relatively light [6]. LHC recently put bounds on the lighter electro-weak chargino-neutralino production cross-sections [7]. If the lighter neutralino states are made of RHsNs which are SM gauge singlets, such production cross-sections are automatically suppressed. The same reason also prompts the non-standard decays of lighter chargions and neutralino which further evade the recent bounds. We consider the $U(1)'$ extended supersymmetric scenario with an right handed sneutrino-superfiled. The $U(1)'$ under question can be $U(1)_{B-L}$ or $U(1)_\chi$ or any other kind of such extensions. For the current study we focus on the $U(1)_\chi$ extended supersymmetric scenarios.

In this work we explore the possible LHC signatures of displaced lepton flavour violating decays of RHsN caused by left-right handed sneutrino mixings. Being SM gauge singlet such RHsNs will mostly come from $\tilde{Z}'$ decay. The pair production cross-section of such $\tilde{Z}'$ is not so promising [6] but its production from SUSY cascade could still be encouraging, in particular from the lighter third generation squark $(\tilde{t}_1, \tilde{b}_1)$ decays. When the heavier RHsN is (Next to next LSP) NNLSP and the lighter one is LSP along with chargino NLSP (Next to LSP), the situation can lead to multi-leptonic final states along with lepton flavour violating displaced charged track.

Among various possibilities of an extra gauge symmetry $U(1)'$ and the presence of the associated right-handed neutrinos [3], we will take the $U(1)_\chi$ model for our explicit analysis as in [6, 8, 9]. The particle content of our $U(1)_\chi$ model is as follows:

$SU(5)$	10_F	$\bar{5}_F$	$1(N)$	5_H	$\bar{5}_H$	$1(X)$	$1(S_1)$	$1(S_2)$
$2\sqrt{10}Q'$	-1	3	-5	2	-2	0	10	-10

(1)

where $SU(5)$ representations and $U(1)'$ charges of the SM fermions $(10_F, \bar{5}_F)$, Higgs bosons $(5_H, \bar{5}_H)$, and additional singlet fields $(N, X, S_{1,2})$ are shown. Here N denotes the right-handed neutrino, X is an additional singlet field fit into the 27 representation of E_6, and we introduced more singlets $S_{1,2}$, vector-like under $U(1)'$, to break $U(1)'$ and generate the Majorana mass term of N [5]. Note that the right-handed neutrinos carry the largest charge under $U(1)'$ and thus the corresponding Z' decays dominantly to right-handed neutrinos. The $U(1)'$ breaking can be generated radiatively in the minimal Higgs sector [5] alternatively the additional singlet field X is neutral under $U(1)'$ so that it can be used to generate a mass to the $U(1)'$ Higgsinos in non-minimal Higgs sectors [10]. Here we do not assume any grand unification theory as the origin of our model and the grand unification structure is just used for a convenient guide to a theoretically consistent model guaranteeing the anomaly free condition.

The gauge invariant superpotential in the seesaw sector

*Electronic address: priyotosh.bandyopadhyay@le.infn.it
†URL: http://www.le.infn.it/~priyo

is given by

$$W_{seesaw} = y_{ij} L_i H_u N_j + \frac{\lambda_{N_i}}{2} S_1 N_i N_i \,, \qquad (2)$$

where L_i and H_u denote the lepton and Higgs doublet superfields, respectively. Given the vacuum expectation values $< S_{1,2} >$, the Z' mass is generated as $2g' \sum_i Q'^2_{S_i} < S_i^2 >$, which gets $\gtrsim 2.5$ TeV bound depending on $U(1)'$ types [11].

After the $U(1)'$ breaking, the right-handed neutrinos (RHNs) obtain the mass $m_{N_i} = \lambda_{N_i} \langle S_1 \rangle$ and induce the seesaw mass for the light neutrinos:

$$\widetilde{m}^{\nu}_{ij} = -y_{ik} y_{jk} \frac{\langle H^0_u \rangle^2}{m_{N_k}} \,. \qquad (3)$$

The same vev also induces the mixing between the RHN N and light neutrino ν and corresponding mixing angle is given by

$$\theta_{N\nu} \simeq \frac{y_\nu v_u}{\sqrt{2} m_N}, \qquad (4)$$

which leads to N decay to SM gauge bosons. The real and imaginary components of the scalar part of N superfiled, $\tilde{N} = (\tilde{N}_1 + \tilde{N}_2)/\sqrt{2}$, get a mass splitting proportional to the lepton number violating (Majorana) mass term m_N[6]. The mixings also occur between the real and imaginary components of the left- and right- sneutrinos proportional to the vev v_u as given by Eq. 5. Here $m_{\tilde{\nu}_{R,I}}$ are masses for the real and imaginary part of the left-handed sneutrino, A_L is the tri-linear soft term corresponding to the leptonic doublet, μ is the up and down type Higgs mixing parameter and $\tan\beta$ is the ratio of their vevs as explained in [6].

$$\theta_{\tilde{N}_{1,2}\tilde{\nu}_{R,I}} = \frac{y_\nu (m_N \pm X_L) v_u}{\sqrt{2}(m^2_{\tilde{N}_{1,2}} - m^2_{\tilde{\nu}_{R,I}})}; \; X_L = A_L \mp \mu \cot\beta \quad (5)$$

These mixing angles can be naturally small due to $y_\nu \sim 10^{-7}$ for $\widetilde{m}^{\nu} = 0.05$ eV and $m_N \sim 100$ GeV and the given the cancelation among the parameters.

An interesting scenario can happen if $\tilde{N}_2$ is NNLSP and $\tilde{N}_1$ is the LSP with a wino like chargino $\tilde{\chi}_1$ as NLSP as shown in Figure 1. If a relatively lighter $\tilde{Z}'$ produced, preferably by SUSY cascade decays, it decays into a RHN and RHsNs ($\tilde{Z}' \to N\tilde{N}_{1,2}$). Such $\tilde{N}_2$ then undergoes a lepton flavour violating decay to an electron and chargino via the mixing $\theta_{\tilde{N}_2,\tilde{\nu}_I}$, with a decay width given in Eq. 6, where V is one of the chargino mixing matrices, g' is the $U(1)'$ gauge coupling and $m_{\tilde{\chi}^\pm_1}$ is the lighter chargino mass. This decay-width can be very small, $\mathcal{O}(10^{-15})$ GeV or less, thus gives rise to displaced charged tracks of length cm to m which can be observed at inner tracker and calorimeters at CMS and ATLAS detectors of the the LHC.

$$\Gamma(\tilde{N}_2 \to e\tilde{\chi}^\pm_1) = \frac{g'^2}{16\pi} |V_{i1}|^2 \theta^2_{\tilde{N}_2,\tilde{\nu}_I} m_{\tilde{N}_2} \left(1 - \frac{m^2_{\tilde{\chi}^\pm_1}}{m^2_{\tilde{N}_2}}\right)^2 \quad (6)$$



FIG. 1: Possible cascade decay of $\tilde{Z}'$ via the mixing of left and right-handed sneutrinos.



FIG. 2: The plausible displaced decay lengths of $\tilde{N}_2$ and $\tilde{\chi}^\pm_1$ in cm in the interested parameter space.

$$\Gamma(\tilde{\chi}^-_1 \to e\tilde{N}_1) = \frac{g'^2}{32\pi} |V_{i1}|^2 \theta^2_{\tilde{N}_1,\tilde{\nu}_R} m_{\tilde{\chi}^\pm_1} \left(1 - \frac{m^2_{\tilde{N}_1}}{m^2_{\tilde{\chi}^\pm_1}}\right)^2 \qquad (7)$$

Such long lived lighter chargino then causes another lepton flavour violating decay to $e^\pm \tilde{N}_1$ for $\Delta m_{\tilde{\chi}^\pm_1,\tilde{N}_1} < m_W$ via snutrino mixing ($\theta_{\tilde{N}_1\tilde{\nu}_R}$). The decay in this case can even be smaller as given in Eq. 7. If the lighter chargino is $\tilde{H}^\pm_u$ type, then $\tilde{N}_2$ decays to $l\tilde{H}^\pm_u$ would be proportional to the y_ν.

Left-right-handed snutrino mixings when go to very small values $\theta_{\tilde{N}_{1,2}\tilde{\nu}_{R,I}} \simeq \mathcal{O}(10^{-8})$, can result in displaced decays for both $\tilde{N}_2$ and $\tilde{\chi}^\pm_1$. Due to the fact $\tilde{N}_2$ is neutral scalar compared to $\tilde{\chi}^\pm_1$ which is a charged fermion, the former decay width is four times more given everything else are the same. For a given m_N and X_L the cancelation happens either for $\theta_{\tilde{N}_1\tilde{\nu}_R}$ or for $\theta_{\tilde{N}_2\tilde{\nu}_I}$.

Figure 2 shows such decay lengths for $\tilde{N}_2$ and $\tilde{\chi}^\pm_1$ in cm for a scanned parameter space, where $100 \le m_N \le 200$, $100 \le m_{\tilde{N}_{1,2}} \le 500$, $205 \le m_{\tilde{\nu}} \le 505$, $m_{\tilde{\chi}^\pm_1} \ge m_{\tilde{N}_1} + 10$ in GeV, $V_{i1} = 1$ and $g' = 0.3$ for $\widetilde{m}^{\nu} = 0.05$ eV. We can see that depending on the parameter space, chargino produced from the first displaced decays of $\tilde{N}_2$ can travel few cm to m before gets another recoil from its decay.

The RHN N can be produced from the Z' decay, $Z' \to NN$, and then can go through a flavour violating decay, $N \to eW$. However, recent bound of $m_{Z'} \gtrsim 2.5$ TeV from LHC [11], reduces such production cross-section drastically[6]. The other source of RHNs could be from the decays of its super-partner $\tilde{Z}'$, $\tilde{Z}' \to N\tilde{N}_{1,2}$. Due



FIG. 3: Electron and muon multiplicity distribution (upper) and their p_T distribution (lower) at the LHC with 14 TeV.

mass	BP1	BP2	BP3
$m_{\tilde{b}_1}$	634.3	656.2	622.7
$m_{\tilde{t}_1}$	561.5	547.4	543.5
$m_{\tilde{Z}'}$	300.0	250.0	300.0
$m_{\tilde{N}_2}$	150.0	140.0	150.0
$m_{\tilde{\chi}_1^\pm}$	121.0	130.0	140.0
$m_{\tilde{N}_1}$	110.0	100.0	110.0
m_N	100.0	100.0	130.0
X_L	89.9	89.0	89.9

TABLE I: The benchmark points for the collider study where $\widetilde{m}^\nu = 0.05$ eV.

mass	BP1	BP2	BP3
$\mathcal{B}(\tilde{b}_1 \to b\tilde{Z}')$	1.00	0.11	1.00
$\mathcal{B}(\tilde{t}_1 \to b\tilde{Z}')$	1.00	0.02	1.00
$\mathcal{B}(\tilde{Z}' \to N\tilde{N}_2)$	0.40	0.28	0.35

TABLE II: Relevant Branching fractions in superysmmetric cascade decay for the benchmark points.

to suppressed pair production of $\tilde{Z}'$, it is encouraged to look for $\tilde{Z}'$ is supersymmetric (SUSY) cascade decays [8]. From [6] (Figure 13), we can see that the right-handed down type squark mostly decays to $\tilde{Z}'$ for the $U(1)_\chi$ model.

Third generation squarks, in particular lighter sbottom ($\tilde{b}_1$) cascade decay to $\tilde{Z}'b$ is our immediate interest. Once produced, the $\tilde{Z}'$ can go through one or two displaced decays as shown in Figure 1. Thus $pp \to \tilde{b}_1\tilde{b}_1^*$ can give rise to $2b+2N+4e+\not{E}_T$, where the two pairs of electron come from the displaced lepton flavour violating decays of $\tilde{N}_2$ and $\tilde{\chi}_1^\pm$ respectively. The RHNs N produced from $\tilde{Z}'$ decay are not displaced but add to the lepton flavour violating decays eW along with $Z\nu$ and $h\nu$ permitted by the phase space. Thus final state can be constituted of $2b+6e+2W+\not{E}_T$, among which four of the electrons are displaced. Now the two Ws can decay both muons and electrons with the same ratios. Checking the final state muon and electron numbers tagged with two b-jets among which 4 electrons are displaced cloud be a clear signature for such cascade decay. Along with $\tilde{b}_1$ contributions, $\tilde{t}_1 \to t\tilde{Z}'$ adds to the final states with more jets or leptons from the top decay.

For the collider study we select few benchamark points given in Table I and we mainly follow Ref [8] for fixing the Higgs sector with decoupled first two generations of squark and gluino. The recent bounds from LHC on Higgs data [1], bounds on third generation SUSY masses [12] and recent bounds on lighter charginos and neutralino production cross-sections [7] are taken into account.

In this collider study we mainly focus on the multileptonic final states tagged with at least two b-jets ($2b+ \geq 6\ell$) along with lepton flavour number, i.e. e and μ. Four of the electrons coming from $\tilde{N}_2$ and $\tilde{\chi}_1^\pm$ are displaced with average decay length of 795.8 and 404.6, 736.9 and 69.9, and 50.5 and 53.6 in cm for BP1, BP2 and BP3 respectively. Table II shows the relevant decay branching fractions involved in SUSY cascade decays. For BP1 and BP2 the RHN is 100 GeV so decays to $W^\pm e^\mp$ and $Z\nu$ with branching fraction of 88%, 12% respectively. However for BP3 as $m_N = 130$ GeV it decays to $h\nu$ as well with branching fraction of $\sim 0.3\%$.

For this purpose we simulate the events coming from $\tilde{t}$s and $\tilde{b}$s via CalcHEP-PYTHIA[13, 14] interface for the final states. The jet formation has been performed using the `Fastjet-3.0.3` [15] with the `CAMBRIDGE AACHEN` algorithm with jet size $R = 0.5$ where the calorimeter coverage is $|\eta| < 4.5$. The threshold p_T cuts on the jets and the leptons are chosen to be 20 GeV and 10 GeV respectively as the leptons in such cascade decays cloud be soft. $\Delta R_{ll} \geq 0.2$ and $\Delta R_{lj} \geq 0.4$ are imposed to isolated the leptons from other leptons and/or jets.

Figure 3 (Upper) shows the electron and muon number multiplicity distribution from $\tilde{b}_1\tilde{b}_1^*$ for BP1 and it is evident that due to lepton flavour violating decays, electron numbers are more than muons. Figure 3 (lower) shows the p_T distributions of such electrons coming from $\tilde{N}_2$ and $\tilde{\chi}_1^\pm$ and they cloud be on softer side due to smaller mass gaps in the corresponding lepton flavour violating decays. Figure 4 shows the transverse decay lengths of $\tilde{N}_2$ and $\tilde{\chi}_1^\pm$ for BP1 at the LHC for ECM of 14 TeV.

Finally in Table III we present the number of events for such final states at an integrated luminosity of 100 fb^{-1} for the center of mass energy of 13 (14) TeV at the LHC.



FIG. 4: Displaced decay length of $\tilde{N}_2$ and $\tilde{\chi}_1^{\pm}$ at the LHC with 14 TeV.

Final states		BP1	BP2	BP3
$\geq 4\ell + 2b$-jets	$\star$	616.7(712.2)	35.0 (49.0)	517.2 (453.2)
	†	542.0(751.3)	10.8(12.0)	624.3 (811.7)
$\geq 6\ell$	$\star$	232.26(215.3)	6.6 (10.7)	114.7(102.0)
	†	485.2(646.6)	6.4(6.9)	411.2(527.1)
$+2b$-jets	$\star$	92.2(79.3)	2.0 (3.5)	38.8(35.4)
	†	65.2(84.8)	0.7(0.8)	43.4(66.1)
$\geq 8e$	$\star$	2.0(1.2)	10^{-2}(0.1)	1.3(1.0)
	†	7.7(7.9)	10^{-2}(10^{-2})	4.1(7.0)
$+2b$-jets	$\star$	1.6(0.7)	10^{-2}(0.1)	0.8(0.6)
	†	2.7(1.5)	10^{-2}(10^{-3})	1.3(3.7)
$\geq 6e + \geq 2\mu$	$\star$	1.8(1.8)	10^{-2}(0.1)	1.3(0.6)
	†	9.5(12.0)	0.1(0.1)	4.7(10.3)
$+2b$-jets	$\star$	1.4(1.2)	10^{-2}(0.1)	0.8(0.5)
	†	2.2(3.1)	10^{-2}(10^{-2})	1.1(2.6))

TABLE III: Final state numbers for the benchmark points at 100 fb^{-1} at the LHC with 13 (14) TeV. $\star$, † correspond to the contributions from $\tilde{b}_1\tilde{b}_1^*$ and $\tilde{t}_1\tilde{t}_1^*$ respectively.

This includes 4ℓ, 6ℓ, $8e$ and $8e+2\mu$ inclusive or/and with two tagged b-jets with single b-jet tagged efficiency of 0.5 [16]. These final states has 4 displaced electrons for the lepton flavour violating signatures. If the lepton flavour is conserved then we expect both displaced electrons and muons in the final sates. The almost background free final states can have some contributions for $4\ell + 2b$-jet final state from SM ZZZ, $t\bar{t}Z$ but without any displaced charged leptons. Thus the signals remain clean with almost no SM backgrounds.

However depending on the benchmark points the cascade decay branching may affect the final state numbers. For example in BP2, though $4\ell + 2b$-jtes and 6ℓ can be observed with few hundreds of fb^{-1} of data, but higher leptonic final states require higher luminosity. BP1 and BP3 cases are favourable points to probe the full decay chain including the electrons coming from the RHNs and electrons and muons from the $W^{\pm}$ bosons, which are not displaced via $8e$ and $6e + 2\mu$ final states. Lighter RHsNs can be probed via their mixings with the left-handed ones through the displaced multi-leptonic final sates at the LHC. Lepton flavour violation if any can be measured by checking the flavour multiplicities in the final states. In case of supersymmetric Type III seesaw a pair of charged RHsN makes the scenario very interesting [17].

PB wants to acknowledge Prof. Eung Jin Chun for the useful discussions.

[1] G. Aad *et al.* [ATLAS and CMS Collaborations], Phys. Rev. Lett. **114**, 191803 (2015) [arXiv:1503.07589 [hep-ex]].

[2] For a review, see, R. N. Mohapatra *et al.*, Rept. Prog. Phys. **70** (2007) 1757 [arXiv:hep-ph/0510213].

[3] P. Langacker, Rev. Mod. Phys. **81** (2009) 1199 [arXiv:0801.1345 [hep-ph]].

[4] For a review, see, G. Jungman, M. Kamionkowski and K. Griest, Phys. Rept. **267** (1996) 195 [arXiv:hep-ph/9506380].

[5] S. Khalil and A. Masiero, Phys. Lett. B **665** (2008) 374 [arXiv:0710.3525 [hep-ph]].

[6] P. Bandyopadhyay, E. J. Chun and J. C. Park, JHEP **1106** (2011) 129 [arXiv:1105.1652 [hep-ph]].

[7] V. Khachatryan *et al.* [CMS Collaboration], Phys. Rev. D **90**, no. 9, 092007 (2014) [arXiv:1409.3168 [hep-ex]]. V. Khachatryan *et al.* [CMS Collaboration], Eur. Phys. J. C **74**, no. 9, 3036 (2014) [arXiv:1405.7570 [hep-ex]].

[8] P. Bandyopadhyay and E. J. Chun, JHEP **1505** (2015) 045 [arXiv:1412.7312 [hep-ph]].

[9] E. J. Chun, Int. J. Mod. Phys. A **30**, no. 15, 1530041 (2015).

[10] U. Ellwanger, C. Hugonie and A. M. Teixeira, Phys. Rept. **496**, 1 (2010) [arXiv:0910.1785 [hep-ph]].

[11] G. Aad *et al.* [ATLAS Collaboration], Phys. Rev. D **90** (2014) 052005 [arXiv:1405.4123 [hep-ex]]. CMS Collaboration, CMS-PAS-EXO-12-061.

[12] G. Aad *et al.* [ATLAS Collaboration], arXiv:1506.08616 [hep-ex]. V. Khachatryan *et al.* [CMS Collaboration], JHEP **1506** (2015) 116 [arXiv:1503.08037 [hep-ex]].

[13] A. Belyaev, N. D. Christensen and A. Pukhov, Comput. Phys. Commun. **184** (2013) 1729 [arXiv:1207.6082 [hep-ph]].

[14] T. Sjostrand, L. Lonnblad and S. Mrenna, [arXiv:hep-ph/0108264].

[15] M. Cacciari, G. P. Salam and G. Soyez, Eur. Phys. J. C **72** (2012) 1896 [arXiv:1111.6097 [hep-ph]].

[16] I. R. Tomalin [CMS Collaboration], J. Phys. Conf. Ser. **110** (2008) 092033.

[17] P. Bandyopadhyay and E. J. Chun, JHEP **1011**, 006 (2010) [arXiv:1007.2281 [hep-ph]].